Baseline Survey for Astragalus barrii Barneby (Barr's Milkvetch) and Physaria didymocarpa var. lanata A. Nels (Woolly Twinpod) in Eastern Big Horn and Southwestern Rosebud Counties, Montana

Prepared for:

The Bureau of Land Management

Prepared by:

Amy Taylor and Richard Caners

Montana Natural Heritage Program Natural Resource Information System Montana State Library

February, 2002



Baseline Survey for Astragalus barrii Barneby (Barris Milkvetch) and Physaria didymocarpa var. lanata A. Nels (Woolly Twinpod) in Eastern Big Horn and Southwestern Rosebud Counties, Montana

Prepared for:
Bureau of Land Management
Miles City Field Office
111 Garryowen Rd. Miles City, MT 59301

Agreement # ESA010009 Task # 5

Prepared by: Amy Taylor, Contract Botanist and Richard Caner, Botany Program Manager







© 2001 Montana Natural Heritage Program

P.O. Box 201800 • 1515 East Sixth Ave • Helena, MT 59620-1800 • 406-444-3009

This document should be cited as follows:

Taylor, A and R. Caners. Baseline Survey for *Astragalus barrii* Barneby (Barrís Milkvetch) and Physaria didymocarpa var. lanata A. Nels. (Woolly Twinpod) In Eastern Big Horn and Southwestern Rosebud Counties, Montana.

Acknowledgements

Many thanks to all the individuals who contributed to this project. A special thanks to Louise de Montigny, Bureau of Land Management, Miles City Field Office, for the provision of logistical support, and to Kent Undlin, Dawn Doran and Jeff Gustad from the field office for their accompaniment in the field. Thanks to Ron Hartman for use of the Rocky Mountain Herbarium for specimen identification, Spring Creek Coal for access to known locations of sensitive plants, and to all landowners who provided access to conduct surveys. Richard Caners, Botany Program Manager, Montana Natural Heritage Program, edited and prepared this document for submission.

Table of Contents

Executive Summary
Introduction
Study Area
Methods
Results4
Discussion and Management Considerations
References8
Figures and Tables
Figure 1. Localities of Astragalus barrii within study area
Figure 2. Localities of <i>Physaria didymocarpa</i> var. <i>lanata</i> within study area10
Table 1. Occurrence records for <i>Astragalus barrii</i> in the project study area11
Table 2. Occurrence records for <i>Physaria didymocarpa</i> var. <i>lanata</i> in the project study area
Appendices
Appendix 1. Pertinent biological information for <i>Astragalus barrii</i> and <i>Physaria didymocarpa</i> var. <i>lanata</i>
Appendix 2. Potential and existing habitat for <i>Astragalus barrii</i> and <i>Physaria didymocarpa</i> var. <i>lanata</i>
Appendix 3. Selected plant and habitat photographs24

Executive Summary

In 2001, The Bureau of Land Management (BLM) Miles City Field Office contracted with the Montana Natural Heritage Program (MTNHP) to conduct baseline surveys for two species of Special Concern, *Astragalus barrii* Barneby (Barr's milkvetch) and *Physaria didymocarpa* var. *lanata* A. Nels. (woolly twinpod), in eastern Big Horn County and southwestern Rosebud County. Both of these species are regional endemics, their overall range being restricted to portions of adjacent Montana, Wyoming and South Dakota.

Both species are known to occur within the study area, but had not been systematically inventoried. Prior to this survey, *P. didymocarpa* var. *lanata* was known in Montana from only one location in Big Horn County, and had been documented from 14 occurrences in north-central Wyoming. *A. barrii* had previously been documented from 33 locations in southeastern Montana, and was known from South Dakota and Wyoming.

Surveys from this study resulted in four new occurrences of *P. didymocarpa* var. *lanata* and one new occurrence of *A. barrii* in the study area. Habitat information including associated plant species, and topographic and edaphic conditions was collected. This information provides additional baseline information for both species in Montana, and serves as a basis for BLM evaluation of the potential impacts of coal bed methane development on these plants and their habitats. The information found in this report may also help clarify the BLM status of these species, and will assist BLM staff with future field identification of both species and their respective habitats.